According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0938-0760. The time required to complete this information collection is estimated to average 52.8 minutes (0.9 minutes per item), including the time to review instructions, search existing data resources, gather the data needed, and complete and review the information collection. If you have comments concerning the accuracy of the time estimate(s) or suggestions for improving this form, please write to: CMS, 7500 Security Boulevard, Attn: PRA Reports Clearance Officer, Mail Stop C4-26-05, Baltimore, Maryland 21244-1850.

Home Health Patient Tracking Sheet

(M0010)	C M S Certification Number:	
(M0014)	Branch State:	
(M0016)	Branch I D Number:	
(M0018)	National Provider Identifier (N P I) for the attending physici	an who has signed the plan of care:
	DK - Unkn	own or Not Available
(M0020)	Patient I D Number:	
(M0030)	Start of Care Date://	
(M0032)	Resumption of Care Date://	□ NA - Not Applicable
	Patient Name:	
(First)	(M I) (Last)	(Suffix)
(M0050)	Patient State of Residence:	
(M0060)	Patient Zip Code:	
(M0063)	Medicare Number: (including suffix)	□ NA - No Medicare
(M0064)	Social Security Number:	☐ UK - Unknown or Not Available
(M0065)	Medicaid Number:	□ NA - No Medicaid
(M0066)	Birth Date:// month / day / year	
(M0069)	Gender:	
	1 - Male	
	2 - Female	
(M0140)	Race/Ethnicity: (Mark all that apply.)	
	1 - American Indian or Alaska Native	
	2 - Asian	
	3 - Black or African-American	
	4 - Hispanic or Latino	
	5 - Native Hawaiian or Pacific Islander	
	6 - White	

(M0150)	Cι	ırr	ent	Payment Sources for Home Care: (Mark all that apply.)
	(0	-	None; no charge for current services
		1	-	Medicare (traditional fee-for-service)
	:	2	-	Medicare (HMO/managed care/Advantage plan)
	;	3	-	Medicaid (traditional fee-for-service)
		4	-	Medicaid (HMO/managed care)
		5	-	Workers' compensation
	(6	-	Title programs (for example: Title III, V, or XX)
		7	-	Other government (for example: TriCare, VA)
	;	8	-	Private insurance
	9	9	-	Private HMO/managed care
	1	0	-	Self-pay
	1	1	-	Other (specify)

☐ UK - Unknown

DRAFT

Outcome and Assessment Information Set Items to be Used at Specific Time Points

Time Point	<u>Items Used</u>			
Start of Care	M0010-M0030, M0040-M0150, M1000-M1036, M1100-M1306, M1308, M1320-M1410, M1600-M2002, M2010, M2020-M2250			
Resumption of Care (after inpatient stay)	M0032, M0080-M0110, M1000-M1036, M1100-M1306, M1308, M1320-M1410, M1600-M2002, M2010, M2020-M2250			
Follow-Up	M0080-M0100, M0110, M1011, M1021-M1030, M1200,			
Recertification (follow-up) assessment Other follow-up assessment	M1242, M1306, M1308, M1322-M1342, M1400, M1610, M1620, M1630, M1810-M1840, M1850, M1860, M2030, M2200			
Transfer to an Inpatient Facility				
Transferred to an inpatient facility—patient not discharged from an agency Transferred to an inpatient facility—patient discharged from agency	M2015, M2300-M2410, M2430, M0903, M0906			
Discharge from Agency — Not to an Inpatient Facility				
Death at home	M0080-M0100, M0903, M0906			
Discharge from agency	M0080-M0100, M1041-M1056, M1230, M1242, M1306-M1342, M1400, M1500-M1620, M1700-M1720, M1740, M1745, M1800-M1890, M2004, M2015-M2030, M2102, M2300-M2420, M0903, M0906			

CLINICAL RECORD ITEMS

(M0080)	Discipline of Person Completing Assessment:
	1-RN ☐ 2-PT ☐ 3-SLP/ST ☐ 4-OT
(M0090)	Date Assessment Completed:// month / day / year
(M0100)	This Assessment is Currently Being Completed for the Following Reason:
	Start/Resumption of Care
	1 - Start of care—further visits planned
	3 - Resumption of care (after inpatient stay)
	Follow-Up
	4 - Recertification (follow-up) reassessment [Go to M0110]
	5 – Other follow-up [Go to M0110]
	Transfer to an Inpatient Facility
	6 - Transferred to an inpatient facility—patient not discharged from agency [Go to M1041]
	7 - Transferred to an inpatient facility—patient discharged from agency [Go to M1041]
	<u>Discharge from Agency — Not to an Inpatient Facility</u>
	8 - Death at home [Go to M0903]
	9 - Discharge from agency [Go to M1041]

(M0102)	Date of Physician-ordered Start of Care (Resumption of Care): If the physician indicated a specific start of care (resumption of care) date when the patient was referred for home health services, record the date specified.
	/ / [Go to M0110, if date entered]
	month / day / year
	NA - No specific SOC date ordered by physician
(M0104)	Date of Referral: Indicate the date that the written or verbal referral for initiation or resumption of care was received by the HHA.
	/ / month / day / year
(M0110)	Episode Timing: Is the Medicare home health payment episode for which this assessment will define a case mix group an early episode or a later episode in the patient's current se uence of ad acent Medicare home health payment episodes?
	1 - Early
	2 - Later
	UK - Unknown
П	NA - Not Applicable: No Medicare case mix group to be defined by this assessment.
	NT HISTORY AND DIAGNOSES From which of the following Inpatient Facilities was the patient discharged within the past 14 days? (Mark all that apply.)
П	1 - Long-term nursing facility (NF)
	NA - Patient was not discharged from an inpatient facility [Go to M1017]
(M1005)	Inpatient Discharge Date (most recent):
	<u>-</u> //
	month / day / year
	UK - Unknown

(M1011) List each Inpatient Diagnosis and ICD-10-C M code at the level of highest specificity for only those conditions treated during an inpatient stay within the last 14 days (no V, W, X, Y, or Z codes or surgical codes):

		-		
				Inpatient Facility Diagnosis ICD-10-C M Code
				·
				·
		NA	-	Not applicable (patient was not discharged from an inpatient facility) [Omit "NA" option on SOC, ROC]
(M1017	-	Med	dical	ses Requiring Medical or Treatment Regimen Change Within Past 14 Days: List the patient's Diagnoses and ICD-10-C M codes at the level of highest specificity for those conditions requiring d medical or treatment regimen within the past 14 days (no V, W, X, Y, or Z codes or surgical codes):
			Cha	nged Medical Regimen Diagnosis ICD-10-C M Code
		a.		
		b.		
		NA	-	Not applicable (no medical or treatment regimen changes within the past 14 days)
(M1018	3)	this pas	pati t 14	ons Prior to Medical or Treatment Regimen Change or Inpatient Stay Within Past 14 Days: If ent experienced an inpatient facility discharge or change in medical or treatment regimen within the days, indicate any conditions that existed <u>prior to</u> the inpatient stay or change in medical or treatment in. (Mark all that apply.)
		1	-	Urinary incontinence
		2	-	Indwelling/suprapubic catheter
		3	-	Intractable pain
		4	-	Impaired decision-making
		5	-	Disruptive or socially inappropriate behavior
		6	-	Memory loss to the extent that supervision required
		7	-	None of the above
		NA	-	No inpatient facility discharge and no change in medical or treatment regimen in past 14 days
		UK	-	Unknown

O١	ΛО	#		
()II	ИΒ	#		

Expiration date TBD

DRAFT

(M1021/1023/1025) Diagnoses, Symptom Control, and Optional Diagnoses: List each diagnosis for which the patient is receiving home care in Column 1, and enter its ICD-10-C M code at the level of highest specificity in Column 2 (diagnosis codes only - no surgical or procedure codes allowed). Diagnoses are listed in the order that best reflects the seriousness of each condition and supports the disciplines and services provided. Rate the degree of symptom control for each condition in Column 2. ICD-10-C M sequencing requirements must be followed if multiple coding is indicated for any diagnoses. If a Z-code is reported in Column 2 in place of a diagnosis that is no longer active (a resolved condition), then optional item M1025 (Optional Diagnoses - Columns 3 and 4) may be completed as per instructions in Appendix D of the OASIS Guidance Manual – see discussion in Column 3 directions below. Diagnoses reported in M1025 will not impact payment but may be used to risk adjust quality measures.

Code each row according to the following directions for each column. Review Appendix D of the OASIS-C1 Guidance Manual for complete directions on correct completion of M1021, M1023 and M1025.

- Column 1: Enter the description of the diagnosis. Sequencing of diagnoses should reflect the seriousness of each condition and support the disciplines and services provided.
- Column 2: Enter the ICD-10-C M code for the condition described in Column 1 no surgical or procedure codes allowed. Codes must be entered at the level of highest specificity and ICD-10-C M coding rules and sequencing requirements must be followed. Note that external cause codes (ICD-10-C M codes beginning with V, W, X, or Y) may not be reported in M1021 (Primary Diagnosis) but may be reported in M1023 (Secondary Diagnoses). Also note that when a Z-code is reported in Column 2, the code for the underlying condition can often be entered in Column 2, as long as it is an active on-going condition impacting home health care.

Rate the degree of symptom control for the condition listed in Column 1. Choose one value that represents the degree of symptom control appropriate for each diagnosis using the following scale:

- 0 Asymptomatic, no treatment needed at this time
- 1 Symptoms well controlled with current therapy
- 2 Symptoms controlled with difficulty, affecting daily functioning; patient needs ongoing monitoring
- 3 Symptoms poorly controlled; patient needs frequent adjustment in treatment and dose monitoring
- 4 Symptoms poorly controlled; history of re-hospitalizations

Note that the rating for symptom control in Column 2 should not be used to determine the sequencing of the diagnoses listed in Column 1. These are separate items and sequencing may not coincide.

Column 3: (OPTIONAL) There is no requirement that HHAs enter a diagnosis code in M1025 (Columns 3 and 4).

Diagnoses reported in M1025 will not impact payment but may be used to risk adjust quality measures

Agencies may choose to report an underlying condition in M1025 (Columns 3 and 4) when:

- a Z-code is reported in Column 2 AND
- the underlying condition for the Z-code in Column 2 is a resolved condition. An example of a resolved condition is uterine cancer that is no longer being treated following a hysterectomy.
- Column 4: (OPTIONAL) If a Z-code is reported in M1021/M1023 (Column 2) and the agency chooses to report a resolved underlying condition that requires multiple diagnosis codes under ICD-10-C M coding guidelines, enter the diagnosis descriptions and the ICD-10-C M codes in the same row in Columns 3 and 4. For example, if the resolved condition is a manifestation code, record the diagnosis description and ICD-10-C M code for the underlying condition in Column 3 of that row and the diagnosis description and ICD-10-C M code for the manifestation in Column 4 of that row. Otherwise, leave Column 4 blank in that row.

(Form on next page)

(M1021 Primary Diagnosi	s & (M1023) Other Diagnoses)	(M1025) Optional Diagnoses (OPTIONAL) (not used for payment)			
Column 1	Column 2	Column 3	Column 4		
Diagnoses (Sequencing of diagnoses should reflect the seriousness of each condition and support the disciplines and services provided)	ICD-10-C M and symptom control rating for each condition. Note that the sequencing of these ratings may not match the sequencing of the diagnoses	May be completed if a Z- code is assigned to Column 2 and the underlying diagnosis is resolved	Complete only if the Optional Diagnosis is a multiple coding situation (for example: a manifestation code)		
Description	ICD-10-C M / Symptom Control Rating	Description/ ICD-10-C M	Description/ ICD-10-C M		
(M1021) Primary Diagnosis	V, W, X, Y codes NOT allowed	V, W, X, Y, Z codes NOT allowed	V, W, X, Y, Z codes NOT allowed		
a	a	a	a		
	□0 □1 □2 □3 □4	()	()		
(M1023) Other Diagnoses	All ICD-10–C M codes allowed	V, W, X, Y, Z codes NOT allowed	V, W, X, Y, Z codes NOT allowed		
b	b	b	b		
	□0 □1 □2 □3 □4	()	()		
C.	c	C	C		
	□0 □1 □2 □3 □4	()	()		
d	d	d	d		
	□0 □1 □2 □3 □4	()	()		
e	e	e	e		
	□0 □1 □2 □3 □4	()	()		
f	f	f	f		
	□0 □1 □2 □3 □4	()	()		

(M1030) Therapies the patient receives at home: (Mark all that apply.)

☐ 1 - Intravenous or infusion therapy (excludes 1P)		1	-	Intravenous or infusion therapy (excludes TI	PN)
---	--	---	---	--	-----

- ☐ 2 Parenteral nutrition (TPN or lipids)
- 3 Enteral nutrition (nasogastric, gastrostomy, jejunostomy, or any other artificial entry into the alimentary canal)
- ☐ 4 None of the above

(M1033)		or Hospitalization: Which of the following signs or symptoms characterize this patient as at risk for all call that apply.)
	· ·	History of falls (2 or more falls - or any fall with an injury - in the past 12 months)
		Unintentional weight loss of a total of 10 pounds or more in the past 12 months
	3 -	Multiple hospitalizations (2 or more) in the past 6 months
	4 -	Multiple emergency department visits (2 or more) in the past 6 months
	5 -	Decline in mental, emotional, or behavioral status in the past 3 months
	6 -	Reported or observed history of difficulty complying with any medical instructions (for example, medications, diet, exercise) in the past 3 months
	7 -	Currently taking six or more medications
	8 -	Currently reports exhaustion
	9 -	Other
	10 -	None of the above
(M1034)	Overa	Il Status: hich description best fits the patient's overall status (Check one)
	0 -	The patient is stable with no heightened risk(s) for serious complications and death (beyond those typical of the patient s age)
	1 -	heightened ris (s) for serious complications and death (beyond those typical of the patient s age).
	2 -	The patient is likely to remain in fragile health and have ongoing high risk(s) of serious complications and death.
	3 -	The patient has serious progressive conditions that could lead to death within a year.
	UK -	The patient's situation is unknown or unclear.
(M1036)	Risk F apply.	actors, either present or past, likely to affect current health status and/or outcome: (Mark all that
	1 -	Smoking
	2 -	Obesity
	3 -	Alcohol dependency
	4 -	Drug dependency
	5 -	None of the above
	UK -	Unknown
(M1041)		nza Vaccine Data Collection Period: Does this episode of care (SOC/ROC to Transfer/Discharge) e any dates on or between October 1 and March 31?
	0 -	No [Go to M1051]
	1 -	Yes
(M1046)	Influer	nza Vaccine Received: Did the patient receive the influen a vaccine for this year's flu season
	1 -	Yes; received from your agency during this episode of care
	2 -	Yes; received from your agency during a prior episode of care
	3 -	Yes; received from another health care provider (for example: physician, pharmacist)
	4 -	No; patient offered and declined
	5 -	No; patient assessed and determined to have medical contraindication(s)
	6 -	No; not indicated - patient does not meet age/condition guidelines for influenza vaccine
	7 -	No; inability to obtain vaccine due to declared shortage
	8 -	No; patient did not receive the vaccine due to reasons other than those listed in responses $4-7$.

0 - No	(M1051) Pneumococcal Vaccine:	Has the patier	nt ever received	the pneumocoo	ccal vaccination (PPV)?		
(M1056) Reason PPV not received: If patient has never received the pneumococcal vaccination (PPV), state reason: 1 - Offered and declined 2 - Assessed and determined to have medical contraindication(s) 3 - Not indicated; patient does not meet age/condition guidelines for PPV 4 - None of the above LiVING ARRANGEMENTS	□ 0 - No							
1 - Offered and declined 2 - Assessed and determined to have medical contraindication(s) 3 - Not indicated; patient does not meet age/condition guidelines for PPV 4 - None of the above 4 - None of the above	☐ 1 - Yes [Go to M150	00 at TRN; Go	to M1230 at DC	;]				
2 - Assessed and determined to have medical contraindication(s) 3 - Not indicated; patient does not meet age/condition guidelines for PPV 4 - None of the above	(M1056) Reason PPV not receive	d: If patient ha	s never receive	d the pneumoco	occal vaccination	(PPV), state reason:		
3 - Not indicated; patient does not meet age/condition guidelines for PPV	☐ 1 - Offered and decl	ined						
LIVING ARRANGEMENTS (M1100) Patient Living Situation: Which of the following best describes the patient's residential circumstance and availability of assistance? (Check one box only.) Availability of Assistance Living Arrangement	☐ 2 - Assessed and de	etermined to ha	ve medical con	traindication(s)				
LIVING ARRANGEMENTS (M1100) Patient Living Situation: Which of the following best describes the patient's residential circumstance and availability of assistance? (Check one box only.) Availability of Assistance Around the Regular Regular Short-term Assistance Around the Clock Regular Nighttime Assistance Around the Clock Around the Clock Around the Clock Around the Clock Around the Short-term Assistance Available Around the Clock Around	•		meet age/condit	tion guidelines f	or PPV			
M1100 Patient Living Situation: Which of the following best describes the patient's residential circumstance and availability of assistance? (Check one box only.) Availability of Assistance	☐ 4 - None of the above	re						
M1100 Patient Living Situation: Which of the following best describes the patient's residential circumstance and availability of assistance? (Check one box only.) Availability of Assistance	LIVING ARRANGEMENTS							
availability of assistance? (Check one box only.) Availability of Assistance Living Arrangement		Which of the f	ollowing best de	escribes the pat	ient's residential	circumstance and		
Living Arrangement				boomboo tho put	ioni o rodiaonilai			
Living Arrangement		Availability of	of Assistance					
b. Patient lives with other person(s) in the home	Living Arrangement				short-term	assistance		
c. Patient lives in congregate situation (for example: assisted living, residential care home) SENSORY STATUS (M1200) Vision (with corrective lenses if the patient usually wears them): 0 - Normal vision: sees adequately in most situations; can see medication labels, newsprint. 1 - Partially impaired: cannot see medication labels or newsprint, but can see obstacles in path, and the surrounding layout; can count fingers at arm's length. 2 - Severely impaired: cannot locate objects without hearing or touching them, or patient nonresponsive. (M1210) Ability to Hear (with hearing aid or hearing appliance if normally used): 0 - Adequate: hears normal conversation without difficulty. 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. 2 - Severely impaired: absence of useful hearing. UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): 0 - Understands: clear comprehension without cues or repetitions. 1 - Usually Understands: understands most conversations, but misses some part/intent of message.	a. Patient lives alone	□ 01	□ 02	□ 03	□ 04	□ 05		
situation (for example: assisted living, residential care home) 11	person(s) in the home	□ 06	□ 07	□ 08	□ 09	□ 10		
 (M1200) Vision (with corrective lenses if the patient usually wears them): ○ - Normal vision: sees adequately in most situations; can see medication labels, newsprint. ○ 1 - Partially impaired: cannot see medication labels or newsprint, but <u>can</u> see obstacles in path, and the surrounding layout; can count fingers at arm's length. ○ 2 - Severely impaired: cannot locate objects without hearing or touching them, or patient nonresponsive. (M1210) Ability to Hear (with hearing aid or hearing appliance if normally used): ○ - Adequate: hears normal conversation without difficulty. ○ 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. ○ 2 - Severely Impaired: absence of useful hearing. ○ UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): ○ - Understands: clear comprehension without cues or repetitions. ○ 1 - Usually Understands: understands most conversations, but misses some part/intent of message. ○ 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 	situation (for example: assisted living, residential	□ 11	□ 12	□ 13	□ 14	□ 15		
 □ 0 - Normal vision: sees adequately in most situations; can see medication labels, newsprint. □ 1 - Partially impaired: cannot see medication labels or newsprint, but <u>can</u> see obstacles in path, and the surrounding layout; can count fingers at arm's length. □ 2 - Severely impaired: cannot locate objects without hearing or touching them, or patient nonresponsive. (M1210) Ability to Hear (with hearing aid or hearing appliance if normally used): □ 0 - Adequate: hears normal conversation without difficulty. □ 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. □ 2 - Severely Impaired: absence of useful hearing. □ UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): □ 0 - Understands: clear comprehension without cues or repetitions. □ 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 	SENSORY STATUS							
 □ 1 - Partially impaired: cannot see medication labels or newsprint, but <u>can</u> see obstacles in path, and the surrounding layout; can count fingers at arm's length. □ 2 - Severely impaired: cannot locate objects without hearing or touching them, or patient nonresponsive. (M1210) Ability to Hear (with hearing aid or hearing appliance if normally used): □ 0 - Adequate: hears normal conversation without difficulty. □ 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. □ 2 - Severely Impaired: absence of useful hearing. □ UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): □ 0 - Understands: clear comprehension without cues or repetitions. □ 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 	(M1200) Vision (with corrective ler	nses if the patie	ent usually wear	s them):				
surrounding layout; can count fingers at arm's length. 2 - Severely impaired: cannot locate objects without hearing or touching them, or patient nonresponsive. (M1210) Ability to Hear (with hearing aid or hearing appliance if normally used): 0 - Adequate: hears normal conversation without difficulty. 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. 2 - Severely Impaired: absence of useful hearing. UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): 0 - Understands: clear comprehension without cues or repetitions. 1 - Usually Understands: understands most conversations, but misses some part/intent of message.	□ 0 - Normal vision: s	ees adequately	in most situation	ons; can see me	edication labels, r	newsprint.		
nonresponsive. (M1210) Ability to Hear (with hearing aid or hearing appliance if normally used): 0 - Adequate: hears normal conversation without difficulty. 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. 2 - Severely Impaired: absence of useful hearing. UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): 0 - Understands: clear comprehension without cues or repetitions. 1 - Usually Understands: understands most conversations, but misses some part/intent of message.	surrounding layo	ut; can count fi	ngers at arm's l	ength.		·		
 (M1210) Ability to Hear (with hearing aid or hearing appliance if normally used): 0 - Adequate: hears normal conversation without difficulty. 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. 2 - Severely Impaired: absence of useful hearing. UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): 0 - Understands: clear comprehension without cues or repetitions. 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 		d: cannot loca	te objects witho	ut hearing or to	uching them, or p	patient		
 □ 0 - Adequate: hears normal conversation without difficulty. □ 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. □ 2 - Severely Impaired: absence of useful hearing. □ UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): □ 0 - Understands: clear comprehension without cues or repetitions. □ 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 	nonresponsive.							
 1 - Mildly to Moderately Impaired: difficulty hearing in some environments or speaker may need to increase volume or speak distinctly. 2 - Severely Impaired: absence of useful hearing. UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): 0 - Understands: clear comprehension without cues or repetitions. 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 	(M1210) Ability to Hear (with hear	ing aid or hear	ing appliance if	normally used):				
increase volume or speak distinctly. 2 - Severely Impaired: absence of useful hearing. UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): 0 - Understands: clear comprehension without cues or repetitions. 1 - Usually Understands: understands most conversations, but misses some part/intent of message.	☐ 0 - Adequate: hears	normal conve	rsation without	difficulty.				
 UK - Unable to assess hearing. (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): □ 0 - Understands: clear comprehension without cues or repetitions. □ 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 	increase volume or speak distinctly.							
 (M1220) Understanding of Verbal Content in patient's own language (with hearing aid or device if used): 0 - Understands: clear comprehension without cues or repetitions. 1 - Usually Understands: understands most conversations, but misses some part/intent of message. 								
 Understands: clear comprehension without cues or repetitions. Usually Understands: understands most conversations, but misses some part/intent of message. 	UK - Unable to assess	s nearing.						
1 - Usually Understands: understands most conversations, but misses some part/intent of message.	(M1220) Understanding of Verba	I Content in pa	atient's own land	guage (with hea	ring aid or device	e if used):		
	☐ 0 - Understands: cle	ear comprehen	sion without cue	es or repetitions				
				ersations, but mi	sses some part/i	ntent of message.		
 2 - Sometimes Understands: understands only basic conversations or simple, direct phrases. Frequently requires cues to understand. 	Frequently require	es cues to und		sic conversation	ns or simple, dire	ct phrases.		
☐ 3 - Rarely/Never Understands.☐ UK - Unable to assess understanding.	•		•					

(M1230)	Spee	ch	and Oral (Verbal) Expression of Language (in patient's own language):
	0	-	Expresses complex ideas, feelings, and needs clearly, completely, and easily in all situations with no observable impairment.
	1	-	Minimal difficulty in expressing ideas and needs (may take extra time; makes occasional errors in word choice, grammar or speech intelligibility; needs minimal prompting or assistance).
	2	-	Expresses simple ideas or needs with moderate difficulty (needs prompting or assistance, errors in word choice, organization or speech intelligibility). Speaks in phrases or short sentences.
	3	-	Has severe difficulty expressing basic ideas or needs and requires maximal assistance or guessing by listener. Speech limited to single words or short phrases.
	4	-	<u>Unable</u> to express basic needs even with maximal prompting or assistance but is not comatose or unresponsive (for example: speech is nonsensical or unintelligible).
	5	-	Patient nonresponsive or unable to speak.
(M1240)			patient had a formal Pain Assessment using a standardized, validated pain assessment tool riate to the patient's ability to communicate the severity of pain)
	0	-	No standardized, validated assessment conducted
	1	-	Yes, and it does not indicate severe pain
	2	-	Yes, and it indicates severe pain
(M1242)	Freq	uei	ncy of Pain Interfering with patient's activity or movement:
	0	-	Patient has no pain
	1	-	Patient has pain that does not interfere with activity or movement
	2	-	Less often than daily
	3	-	Daily, but not constantly
	4	-	All of the time
INTEGL	JME	١T	ARY STATUS
(M1300)	Pres	sur	re Ulcer Assessment: Was this patient assessed for Risk of Developing Pressure Ulcers?
	0	-	No assessment conducted [Go to M1306]
	1	-	Yes, based on an evaluation of clinical factors (for example: mobility, incontinence, nutrition) without use of standardized tool
	2	-	Yes, using a standardized, validated tool (for example: Braden Scale, Norton Scale)
(M1302)	Does	thi	s patient have a Risk of Developing Pressure Ulcers?
	0	-	No
	1	-	Yes
(M1306)			is patient have at least one Unhealed Pressure Ulcer at Stage II or Higher or designated as eable"? (EXCLUDES Stage I pressure ulcers and healed Stage II ulcers)
	0	-	No [<i>Go to M1322</i>]
	1	-	Yes
(M1307)			dest Non-epithelialized Stage II Pressure Ulcer that is present at discharge:
	0	-	Was present at the most recent SOC/ROC assessment
	2		Developed since the most recent SOC/ROC assessment. Record date pressure ulcer first identified:
	NΙΛ		nonth / day / year
\sqcup	NA	-	No non-epithelialized Stage II pressure ulcers are present at discharge

(M1308) Current Number of Unhealed Pressure Ulcers at Each Stage or Unstageable:

(Enter 0 if none; EXCLUDES Stage I pressure ulcers and healed Stage II ulcers)

Stage Descriptions—unhealed pressure ulcers				
a.	Stage II: Partial thickness loss of dermis presenting as a shallow open ulcer with red pink wound bed, without slough. May also present as an intact or open/ruptured serum-filled blister.			
b.	Stage III: Full thickness tissue loss. Subcutaneous fat may be visible but bone, tendon, or muscles are not exposed. Slough may be present but does not obscure the depth of tissue loss. May include undermining and tunneling.			
C.	Stage IV: Full thickness tissue loss with visible bone, tendon, or muscle. Slough or eschar may be present on some parts of the wound bed. Often includes undermining and tunneling.			
d.1	Unstageable: Known or likely but unstageable due to non-removable dressing or device			
d.2	Unstageable: Known or likely but unstageable due to coverage of wound bed by slough and/or eschar.			
d.3	Unstageable: Suspected deep tissue injury in evolution.			

(M1309) Worsening in Pressure Ulcer Status since SOC/ROC: Indicate the number of current pressure ulcers that were not present or were at a lesser stage at the most recent SOC/ROC: (Enter 0 if none)

Stage	Number of current pressure ulcers that were NOT PRESENT or were at a LESSER STAGE at most recent SOC/ROC
a. Stage II	
b. Stage III	
c. Stage IV	_

0	-	Newly epithelialized
1	-	Fully granulating
2	-	Early/partial granulation
3	-	Not healing
NA	-	No observable pressure ulcer

(M1322) Current Number of Stage I Pressure Ulcers: Intact skin with non-blanchable redness of a localized area usually over a bony prominence. The area may be painful, firm, soft, warmer, or cooler as compared to adjacent tissue.

□ 0	□ 1	□ 2	□ 3	☐ 4 or more
-----	-----	-----	-----	-------------

(M1324) St	age of Most Problematic Unhealed (Observable) Pressure Ulcer:
	1 - Stage I
	2 - Stage II
	3 - Stage III
	4 - Stage IV
□ N/	A - No observable pressure ulcer or unhealed pressure ulcer
(M1330) Do	es this patient have a Stasis Ulcer ?
	O - No [<i>Go to M1340</i>]
	1 - Yes, patient has BOTH observable and unobservable stasis ulcers
	2 - Yes, patient has observable stasis ulcers ONLY
	 Yes, patient has unobservable stasis ulcers ONLY (known but not observable due to non-removable dressing) [Go to M1340]
(M1332) Cu	rrent Number of (Observable) Stasis Ulcer(s):
	1 - One
	2 - Two
	3 - Three
	4 - Four or more
(M1334) St	atus of Most Problematic (Observable) Stasis Ulcer:
	1 - Fully granulating
	2 - Early/partial granulation
	3 - Not healing
(M1340) Do	es this patient have a Surgical Wound?
	- No [At SOC/ROC, go to M1350; At TRN/DC, go to M1400]
	1 - Yes, patient has at least one (observable) surgical wound
	 Surgical wound known but not observable due to non-removable dressing [At SOC/ROC, go to M1350; At TRN/DC, go to M1400]
(M1342) St	atus of Most Problematic (Observable) Surgical Wound:
	O - Newly epithelialized
	1 - Fully granulating
	2 - Early/partial granulation
	3 - Not healing
	nes this patient have a Skin Lesion or Open Wound (excluding bowel ostomy), other than those described ove, that is receiving intervention by the home health agency?
) - No
	1 - Yes

OMB #_____ Expiration date TBD

DRAFT

RESPIRATORY STATUS

(M1400)	Wh	en i	s the patient dyspneic or noticeably Short of Breath?
	0	-	Patient is not short of breath
	1	-	When walking more than 20 feet, climbing stairs
	2	-	With moderate exertion (for example: while dressing, using commode or bedpan, walking distances less than 20 feet)
	3	-	
	4	-	At rest (during day or night)
(M1410)	Res	spira	atory Treatments utilized at home: (Mark all that apply.)
	1	_	Oxygen (intermittent or continuous)
	2	_	Ventilator (continually or at night)
	_	_	
		_	
CARDI	AC	ST	ATUS
	Syr syr	npto	oms in Heart Failure Patients: If patient has been diagnosed with heart failure, did the patient exhibit oms indicated by clinical heart failure guidelines (including dyspnea, orthopnea, edema, or weight gain) time of or at any time since the previous OASIS assessment?
	0	-	No [Go to M2004 at TRN; Go to M1600 at DC]
	1	-	Yes
	2	-	Not assessed [Go to M2004 at TRN; Go to M1600 at DC]
	NA	-	Patient does not have diagnosis of heart failure [Go to M2004 at TRN; Go to M1600 at DC]
(M1510)	ind	icati	Failure Follow-up: If patient has been diagnosed with heart failure and has exhibited symptoms ve of heart failure at the time of or at any time since the previous OASIS assessment, what action(s) ave) been taken to respond? (Mark all that apply.)
	0	-	No action taken
	1	-	Patient's physician (or other primary care practitioner) contacted the same day
	2	-	Patient advised to get emergency treatment (for example: call 911 or go to emergency room)
	3	-	Implemented physician-ordered patient-specific established parameters for treatment
	4	-	Patient education or other clinical interventions
	5	-	Obtained change in care plan orders (for example: increased monitoring by agency, change in visit frequency, telehealth)
			07.471.0
			I STATUS
(M1600)	Has	s this	s patient been treated for a Urinary Tract Infection in the past 14 days?
	0	-	No
	1	-	Yes
	NA	-	Patient on prophylactic treatment
	UK	-	Unknown [Omit "UK" option on DC]

(M161	0)	Urin	ary	Incontinence or Urinary Catheter Presence:
		0	-	No incontinence or catheter (includes anuria or ostomy for urinary drainage) [Go to M1620]
		1	-	Patient is incontinent
		2	-	Patient requires a urinary catheter (specifically: external, indwelling, intermittent, or suprapubic) [Go to M1620]
(M161	5)	Whe	n d	oes Urinary Incontinence occur?
		0	-	Timed-voiding defers incontinence
		1	-	Occasional stress incontinence
		2	-	During the night only
		3	-	During the day only
		4	-	During the day and night
(M162	0) B	owe	el In	continence Frequency:
		0	-	Very rarely or never has bowel incontinence
		1	-	Less than once weekly
		2	-	One to three times weekly
		3	-	Four to six times weekly
		4	-	On a daily basis
		5	-	More often than once daily
		NA	-	Patient has ostomy for bowel elimination
		UK	-	Unknown [Omit "UK" option on FU, DC]
(M163	-		ays	r for Bowel Elimination: Does this patient have an ostomy for bowel elimination that (within the last) a) was related to an inpatient facility stay, <u>or</u> b) necessitated a change in medical or treatment ?
		0	-	Patient does not have an ostomy for bowel elimination.
		1	-	Patient's ostomy was <u>not</u> related to an inpatient stay and did <u>not</u> necessitate change in medical or treatment regimen.
		2	-	The ostomy was related to an inpatient stay or did necessitate change in medical or treatment regimen.
<u>NEUI</u>	₹0/	/EM	01	TIONAL/BEHAVIORAL STATUS
(M170				ve Functioning: Patient's current (day of assessment) level of alertness, orientation, hension, concentration, and immediate memory for simple commands.
		0	-	Alert/oriented, able to focus and shift attention, comprehends and recalls task directions independently.
		1	-	Requires prompting (cuing, repetition, reminders) only under stressful or unfamiliar conditions.
		2	-	Requires assistance and some direction in specific situations (for example: on all tasks involving shifting of attention) or consistently requires low stimulus environment due to distractibility.
		3	-	Requires considerable assistance in routine situations. Is not alert and oriented or is unable to shift attention and recall directions more than half the time.
		4	-	Totally dependent due to disturbances such as constant disorientation, coma, persistent vegetative state, or delirium.

(M1710)) \	Nh€	n C	Confused (Reported or 0	Observed Wi	thin the Last	14 Days):		
1		0	-	Never					
1		1	-	In new or complex situa	tions only				
1		2	-	On awakening or at nigh	nt only				
I		3	-	During the day and ever	ning, but not c	constantly			
I		4	-	Constantly					
1	<u> </u>	NΑ	-	Patient nonresponsive					
(M1720)) \	Νhe	n A	Anxious (Reported or O	bserved With	nin the Last 1	4 Days):		
		0	-	None of the time					
		1	-	Less often than daily					
		2	-	Daily, but not constantly					
1		3	-	All of the time					
1	<u> </u>	NΑ	-	Patient nonresponsive					
		depi 0	ess	sion Screening: Has the sion screening tool? No Yes, patient was screen patient: Over the last to problems)	ed using the I	PHQ-2©* scal	e. (Instructions	for this two-q	uestion tool: Ask
				PHQ-2©*	Not at all 0 - 1 day	Several days 2 - 6 days	More than half of the days 7 – 11 days	Nearly every day 12 – 14 days	N/A Unable to respond
Ī	a)			interest or pleasure in	0	<u></u> 1	□2	□3	□NA
-	b)	Fe	elir	things ng down, depressed, or ess?	□0	1	□2	□3	□NA
1	□ □ * <i>Co</i>	3		Yes, patient was screen meets criteria for further Yes, patient was screen not meet criteria for furth Pfizer Inc. All rights res	evaluation fo ed with a diffe ner evaluation	or depression. erent standard n for depressio	lized, validated a		·
M1740) (og or O	niti bse	ve, behavioral, and psy erved): (Mark all that ap	chiatric sym _l oply.)	ptoms that ar	e demonstrated	at least once	a week (Report
1		1	-	Memory deficit: failure thours, significant memory				ty to recall eve	ents of past 24
-		2	-	Impaired decision-makir activities, jeopardizes sa	afety through	actions			
l		3	-	Verbal disruption: yellin	-		-		
I		4	-	Physical aggression: agobjects, punches, dange	erous maneuv	ers with whee	elchair or other o	objects)	
l		5	-	Disruptive, infantile, or s		•	ior (excludes v	erbal actions)	
I		6	-	Delusional, hallucinatory	, or paranoid	behavior			
		7	-	None of the above beha	viors demons	strated			

(M1745)			ncy of Disruptive Behavior Symptoms (Reported or Observed): Any physical, verbal, or other ve/dangerous symptoms that are injurious to self or others or jeopardize personal safety.
	0	-	Never
	1	-	Less than once a month
	2	-	Once a month
	3	-	Several times each month
	4	-	Several times a week
	5	-	At least daily
(M1750)	ls th	nis p	atient receiving Psychiatric Nursing Services at home provided by a qualified psychiatric nurse?
	0	-	No
	1	-	Yes
ADL/IA	DLs	<u>i</u>	
(M1800)			ng: Current ability to tend safely to personal hygiene needs (specifically: washing face and hands, e, shaving or make up, teeth or denture care, or fingernail care).
	0	-	Able to groom self unaided, with or without the use of assistive devices or adapted methods.
	1	-	Grooming utensils must be placed within reach before able to complete grooming activities.
	2	-	Someone must assist the patient to groom self.
	3	-	Patient depends entirely upon someone else for grooming needs.
(M1810)			Ability to Dress <u>Upper</u> Body safely (with or without dressing aids) including undergarments, s, front-opening shirts and blouses, managing zippers, buttons, and snaps:
	0	-	Able to get clothes out of closets and drawers, put them on and remove them from the upper body without assistance.
	1	-	Able to dress upper body without assistance if clothing is laid out or handed to the patient.
	2	-	Someone must help the patient put on upper body clothing.
	3	-	Patient depends entirely upon another person to dress the upper body.
(M1820)			Ability to Dress <u>Lower</u> Body safely (with or without dressing aids) including undergarments, slacks, r nylons, shoes:
	0	-	Able to obtain, put on, and remove clothing and shoes without assistance.
	1	-	Able to dress lower body without assistance if clothing and shoes are laid out or handed to the patient.
	2	-	Someone must help the patient put on undergarments, slacks, socks or nylons, and shoes.
	3	-	Patient depends entirely upon another person to dress lower body.

(M1830				: Current ability to wash entire body safely. <u>Excludes</u> grooming (washing face, washing hands, impooing hair).
[0	-	Able to bathe self in shower or tub independently, including getting in and out of tub/shower.
[1	-	With the use of devices, is able to bathe self in shower or tub independently, including getting in and out of the tub/shower.
[2	-	Able to bathe in shower or tub with the intermittent assistance of another person: (a) for intermittent supervision or encouragement or reminders, <u>OR</u> (b) to get in and out of the shower or tub, <u>OR</u> (c) for washing difficult to reach areas.
[3	-	Able to participate in bathing self in shower or tub, <u>but</u> requires presence of another person throughout the bath for assistance or supervision.
[4	-	Unable to use the shower or tub, but able to bathe self independently with or without the use of devices at the sink, in chair, or on commode.
[5	-	Unable to use the shower or tub, but able to participate in bathing self in bed, at the sink, in bedside chair, or on commode, with the assistance or supervision of another person.
[6	-	Unable to participate effectively in bathing and is bathed totally by another person.
(M1840				ransferring: Current ability to get to and from the toilet or bedside commode safely and transfer on oilet/commode.
[0	-	Able to get to and from the toilet and transfer independently with or without a device.
[1	-	When reminded, assisted, or supervised by another person, able to get to and from the toilet and transfer.
[2	-	<u>Unable</u> to get to and from the toilet but is able to use a bedside commode (with or without assistance).
[3		<u>Unable</u> to get to and from the toilet or bedside commode but is able to use a bedpan/urinal independently.
L	┙	4	-	Is totally dependent in toileting.
(M1845		pads	s be	g Hygiene: Current ability to maintain perineal hygiene safely, adjust clothes and/or incontinence fore and after using toilet, commode, bedpan, urinal. If managing ostomy, includes cleaning area stoma, but not managing equipment.
[0	-	Able to manage toileting hygiene and clothing management without assistance.
[1		Able to manage toileting hygiene and clothing management without assistance if supplies/implements are laid out for the patient.
[2	-	Someone must help the patient to maintain toileting hygiene and/or adjust clothing.
[3	-	Patient depends entirely upon another person to maintain toileting hygiene.
(M1850				rring: Current ability to move safely from bed to chair, or ability to turn and position self in bed if s bedfast.
[0	-	Able to independently transfer.
[1	-	Able to transfer with minimal human assistance or with use of an assistive device.
[2	-	Able to bear weight and pivot during the transfer process but unable to transfer self.
[3	-	Unable to transfer self and is unable to bear weight or pivot when transferred by another person.
[4	-	Bedfast, unable to transfer but is able to turn and position self in bed.
[5	-	Bedfast, unable to transfer and is unable to turn and position self.

(M1860)			ition/Locomotion: Current ability to walk safely, once in a standing position, or use a wheelchair, a seated position, on a variety of surfaces.
	0	-	Able to independently walk on even and uneven surfaces and negotiate stairs with or without railings (specifically: needs no human assistance or assistive device).
	1	-	With the use of a one-handed device (for example: cane, single crutch, hemi-walker), able to independently walk on even and uneven surfaces and negotiate stairs with or without railings.
	2	-	Requires use of a two-handed device (for example: walker or crutches) to walk alone on a level
			surface and/or requires human supervision or assistance to negotiate stairs or steps or uneven surfaces.
	3	-	Able to walk only with the supervision or assistance of another person at all times.
	4	-	Chairfast, <u>unable</u> to ambulate but is able to wheel self independently.
	5	-	Chairfast, unable to ambulate and is <u>unable</u> to wheel self.
	6	-	Bedfast, unable to ambulate or be up in a chair.
(M1870)			g or Eating: Current ability to feed self meals and snacks safely. Note: This refers only to the of eating, chewing, and swallowing, not preparing the food to be eaten.
	0	-	Able to independently feed self.
	1	-	Able to feed self independently but requires:
			 (a) meal set-up; <u>OR</u> (b) intermittent assistance or supervision from another person; <u>OR</u> (c) a liquid, pureed or ground meat diet.
	2	-	<u>Unable</u> to feed self and must be assisted or supervised throughout the meal/snack.
	3	-	Able to take in nutrients orally <u>and</u> receives supplemental nutrients through a nasogastric tube or gastrostomy.
	4	-	<u>Unable</u> to take in nutrients orally and is fed nutrients through a nasogastric tube or gastrostomy.
	5	-	Unable to take in nutrients orally or by tube feeding.
(M1880)	Curre		Ability to Plan and Prepare Light Meals (for example: cereal, sandwich) or reheat delivered meals
	0	-	(a) Able to independently plan and prepare all light meals for self or reheat delivered meals; <u>OR</u>(b) Is physically, cognitively, and mentally able to prepare light meals on a regular basis but has not routinely performed light meal preparation in the past (specifically: prior to this home care admission).
	1	-	<u>Unable</u> to prepare light meals on a regular basis due to physical, cognitive, or mental limitations.
	2	-	Unable to prepare any light meals or reheat any delivered meals.
(M1890)			to Use Telephone: Current ability to answer the phone safely, including dialing numbers, and ely using the telephone to communicate.
	0	-	Able to dial numbers and answer calls appropriately and as desired.
	1	-	Able to use a specially adapted telephone (for example: large numbers on the dial, teletype phone for the deaf) and call essential numbers.
	2	-	$\label{thm:conversation} \mbox{Able to answer the telephone and carry on a normal conversation but has difficulty with placing calls.}$
	3	-	lem:lem:lem:lem:lem:lem:lem:lem:lem:lem:
	4	-	<u>Unable</u> to answer the telephone at all but can listen if assisted with equipment.
	5	-	Totally unable to use the telephone.
	NA	-	Patient does not have a telephone.

(M1900) Prior Functioning ADL/IADL: Indicate the patient's usual ability with everyday activities prior to this current illness, exacerbation, or injury. Check only <u>one</u> box in each row.

	Functional Area	Independent	Needed Some Help	Dependent
a.	Self-Care (specifically: grooming, dressing, bathing, and toileting hygiene)	□0	□ 1	□2
b.	Ambulation	□0	□1	□2
c.	Transfer	□0	□1	□2
d.	Household tasks (specifically: light meal preparation, laundry, shopping, and phone use)	□0	□1	□2

(M1910) Has this patient had a multi-factor Falls Risk Assessment using a standardized, validated assessment tool?

[0	-	No.
		1	-	Yes, and it indicated no, low, or minimal risk for falls.
[2	-	Yes, and it indicated more than a minimal risk for falls.
<u>MEDI</u>	C	ATI	10	<u>us</u>
(M2000	•	med	licat	egimen Review: Does a complete drug regimen review indicate potential clinically significant ion issues (for example: adverse drug reactions, ineffective drug therapy, significant side effects, drug ions, duplicate therapy, omissions, dosage errors, or noncompliance [non-adherence])?
		0	-	Not assessed/reviewed [Go to M2010]
		1	-	No problems found during review [Go to M2010]
[2	-	Problems found during review
[NA	-	Patient is not taking any medications [Go to M2040]
(M2002	•			tion Follow-up: Was a physician or the physician-designee contacted within one calendar day to clinically significant medication issues, including reconciliation?
		0	-	No
[1	-	Yes
(M2004	•	time	sin	tion Intervention: If there were any clinically significant medication issues at the time of, or at any ce the previous OASIS assessment, was a physician or the physician-designee contacted within one or day to resolve any identified clinically significant medication issues, including reconciliation?
		0	-	No
[1	-	Yes
[NA	-	No clinically significant medication issues identified at the time of or at any time since the previous OASIS assessment

(M2010	O10) Patient/Caregiver High-Risk Drug Education: Has the patient/caregiver received instruction on specific precautions for all high-risk medications (such as hypoglycemics, anticoagulants, etc.) and how and very problems that may occur?									
		0	-	No						
		1	-	Yes						
		NA	-	Patient not taking a precautions assoc		s OR patient/caregiver risk medications	er fully knowledgeat	ole about special		
(M2015		asse effe	essn ctive	nent, was the patier	nt/caregiver instructory, adverse drug r	ntion: At the time of, coted by agency staff of eactions, and significations.	or other health care	provider to monitor the		
		0	-	No						
		1	-	Yes						
		NA	-	Patient not taking	any drugs					
(M2020		and	safe	ely, including admin	istration of the cor	s current ability to pre- rect dosage at the ap is refers to ability, n	propriate times/inte			
		0	-	Able to independe	ntly take the corre	ct oral medication(s)	and proper dosage	(s) at the correct times.		
ĺ		1	-	Able to take medic	ation(s) at the cor	rect times if:				
		(a) individual dosages are prepared in advance by another person; <u>OR</u>(b) another person develops a drug diary or chart.								
		2	-	Able to take medic appropriate times	ation(s) at the cor	rect times if given rer	minders by another	person at the		
		3	-	<u>Unable</u> to take me	dication unless ad	lministered by anothe	er person.			
		NA	-	No oral medication	s prescribed.					
injectal	ole	med	icat			atient's current ability ninistration of correct		e <u>all</u> prescribed opriate times/intervals.		
		0 - Able to independently take the correct medication(s) and proper dosage(s) at the correct times.								
		1 - Able to take injectable medication(s) at the correct times if:								
		(a) individual syringes are prepared in advance by another person; <u>OR</u>(b) another person develops a drug diary or chart.								
		2	 Able to take medication(s) at the correct times if given reminders by another person based on the frequency of the injection 							
		3	-	Unable to take inje	ctable medication	unless administered	by another person.			
		NA	-	No injectable med	cations prescribed	d.				
(M2040						e patient s usual abili				
		F	unc	ctional Area	Independent	Needed Some Help	Dependent	Not Applicable		
	a.	Ora	l me	edications	□0	□1	□2	□NA		
	b.	Inje	ctab	le medications	□0	□1	□2	□NA		

CARE MANAGEMENT

(M2102) Types and Sources of Assistance: Determine the ability and willingness of non-agency caregivers (such as family members, friends, or privately paid caregivers) to provide assistance for the following activities, if assistance is needed. EXCLUDES all care by your agency staff. (Check only **one** box in each row.)

Type of Assistance	No assistance needed – patient is independent or does not have needs in this area	Non-agency caregiver(s) currently provide assistance	Non-agency caregiver(s) need training/ supportive services to provide assistance	Non-agency caregiver(s) are not likely to provide assistance OR it is unclear if they will provide assistance	Assistance needed, but no non-agency caregiver(s) available	
a. ADL assistance (for example: transfer/ ambulation, bathing, dressing, toileting, eating/feeding)	□0	□1	□2	□3	□4	
b. IADL assistance (for example: meals, housekeeping, laundry, telephone, shopping, finances)	□0	□1	□2	□3	□4	
c. Medication administration (for example: oral, inhaled or injectable)	□0	□1	□2	□3	□4	
d. Medical procedures/ treatments (for example: changing wound dressing)	□0	□ 1	□2	□3	□4	
e. Management of Equipment (for example: oxygen, IV/infusion equip- ment, enteral/ parenteral nutrition, ventilator therapy equipment or supplies)	□0	□1	□2	□3	□ 4	
f. Supervision and safety (for example: due to cognitive impairment)	□0	□1	□2	□3	<u></u> 4	
g. Advocacy or facilitation of patient's participation in appropriate medical care (for example: transportation to or from appointments)	□0	□1	□2	□3	□4	

(M2110) How Often does the patient receive AD agency staff)?	L or IAD	L assista	ance from	n any caregiver(s) (other than home health			
☐ 1 - At least daily							
☐ 2 - Three or more times per week							
☐ 3 - One to two times per week							
4 - Received, but less often than w	veekly						
5 - No assistance received	· comy						
☐ UK - Unknown [Omit "UK" option o	on DC1						
OK - OHKHOWIT CHILL OK OPTION	on DC]						
THERAPY NEED AND PLAN OF CARE							
(M2200) Therapy Need: In the home health plat will define a case mix group, what is the physical, occupational, and speech-land therapy visits indicated.)	indicate	d need fo	r therapy	visits (total of reasonable and necessary			
() Number of therapy visits indica combined).							
☐ NA - Not Applicable: No case mix g	roup defi	ned by th	is assess	ment.			
(M2250) Plan of Care Synopsis: (Check only on include the following:	one box ir	n each ro	w.) Does	the physician-ordered plan of care			
Plan / Intervention	No	Yes	Not Ap	olicable			
Patient-specific parameters for notifying physician of changes in vital signs or other clinical findings	□0	□1	□NA	Physician has chosen not to establish patient-specific parameters for this patient. Agency will use standardized clinical guidelines accessible for all care providers to reference.			
 Diabetic foot care including monitoring for the presence of skin lesions on the lower extremities and patient/caregiver education on proper foot care 	□0	□1	□NA	Patient is not diabetic or is missing lower legs due to congenital or acquired condition (bilateral amputee).			
c. Falls prevention interventions	□0	□1	□NA	Falls risk assessment indicates patient has no, low, or minimal risk for falls.			
d. Depression intervention(s) such as medication, referral for other treatment, or a monitoring plan for current treatment	<u></u> 0	<u></u> 1	□NA	Patient has no diagnosis of depression AND depression screening indicates patient has: 1) no symptoms of depression; or 2) has some symptoms of depression but does not meet criteria for further evaluation of depression based on screening tool used.			
e. Intervention(s) to monitor and mitigate pain	□0	□1	□NA	Pain assessment indicates patient has no pain.			
f. Intervention(s) to prevent pressure ulcers	□0	<u></u> 1	□NA	Pressure ulcer risk assessment (clinical or formal) indicates patient is not at risk of developing pressure ulcers.			
g. Pressure ulcer treatment based on principles of moist wound healing OR order for treatment based on moist wound healing has been requested from physician	□ 0	<u></u> 1	□NA	Patient has no pressure ulcers OR has no pressure ulcers for which moist wound healing is indicated.			

DRAFT

EMERGENT CARE

(M2300)		_	ent Care: At the time of or at any time since the previous OASIS assessment has the patient utilized tal emergency department (includes holding/observation status)?
	0	-	No [<i>Go to M2400</i>]
	1	-	Yes, used hospital emergency department WITHOUT hospital admission
	2	-	Yes, used hospital emergency department WITH hospital admission
	UK	-	Unknown [Go to M2400]
(M2310)			for Emergent Care: For what reason(s) did the patient seek and/or receive emergent care (with or hospitalization)? (Mark all that apply.)
	1	-	Improper medication administration, adverse drug reactions, medication side effects, toxicity, anaphylaxis
	2	-	Injury caused by fall
	3	-	Respiratory infection (for example: pneumonia, bronchitis)
	4	-	Other respiratory problem
	5	-	Heart failure (for example: fluid overload)
	6	-	Cardiac dysrhythmia (irregular heartbeat)
	7	-	Myocardial infarction or chest pain
	8	-	Other heart disease
	9	-	Stroke (CVA) or TIA
	10	-	Hypo/Hyperglycemia, diabetes out of control
	11	-	GI bleeding, obstruction, constipation, impaction
	12	-	Dehydration, malnutrition
	13	-	Urinary tract infection
	14	-	IV catheter-related infection or complication
	15	-	Wound infection or deterioration
	16	-	Uncontrolled pain
	17	-	Acute mental/behavioral health problem
	18	-	Deep vein thrombosis, pulmonary embolus
	19	-	Other than above reasons
П	UK	-	Reason unknown

DRAFT

<u>DATA ITEMS COLLECTED AT INPATIENT FACILITY ADMISSION OR AGENCY DISCHARGE ONLY</u>

(M2400) Intervention Synopsis: (Check only <u>one</u> box in each row.) At the time of or at any time since the previous OASIS assessment, were the following interventions BOTH included in the physician-ordered plan of care AND implemented?

	Plan / Intervention	No	Yes	Not App	olicable
a.	Diabetic foot care including monitoring for the presence of skin lesions on the lower extremities and patient/caregiver education on proper foot care	□0	<u></u> 1	□NA	Patient is not diabetic or is missing lower legs due to congenital or acquired condition (bilateral amputee).
b.	Falls prevention interventions	□0	<u></u> 1	□NA	Every standardized, validated multi- factor fall risk assessment conducted at or since the last OASIS assessment indicates the patient has no, low or minimal risk for falls.
C.	Depression intervention(s) such as medication, referral for other treatment, or a monitoring plan for current treatment	O	_1	□NA	Patient has no diagnosis of depression AND every standardized, validated depression screening conducted at or since the last OASIS assessment indicates the patient has: 1) no symptoms of depression; or 2) has some symptoms of depression but does not meet criteria for further evaluation of depression based on screening tool used.
d.	Intervention(s) to monitor and mitigate pain	□0	<u></u> 1	□NA	Every standardized, validated pain assessment conducted at or since the last OASIS assessment indicates the patient has no pain.
e.	Intervention(s) to prevent pressure ulcers	□0	<u></u> 1	□NA	Every standardized, validated pressure ulcer risk assessment conducted at or since the last OASIS assessment indicates the patient is not at risk of developing pressure ulcers.
f.	Pressure ulcer treatment based on principles of moist wound healing	□0	□ 1	□NA	Patient has no pressure ulcers OR has no pressure ulcers for which moist wound healing is indicated.
(M:	2410) To which Inpatient Facility has the	patient b	een admi	tted?	
`	☐ 1 - Hospital [<i>Go to M2430</i>]				
	☐ 2 - Rehabilitation facility [Go	to M0903	³]		
	☐ 3 - Nursing home [Go to M0	903]			
	☐ 4 - Hospice [<i>Go to M0903</i>]				
	☐ NA - No inpatient facility admiss	ion [Omit	"NA" op	otion on T	RN]
(M:	2420) Discharge Disposition: Where is answer.)	the patier	nt after dis	scharge fr	om your agency? (Choose only one
	☐ 1 - Patient remained in the co	mmunity (without fo	ormal assi	stive services)
	☐ 2 - Patient remained in the co	mmunity (with form	al assistiv	re services)
	☐ 3 - Patient transferred to a not	n-institutio	nal hospi	ice	
			a geogra	phic locat	ion not served by this agency
	☐ UK - Other unknown [Go to Me	<i>0903</i>]			

(M2430)	Rea		for Hospitalization: For what reason(s) did the patient require hospitalization? (Mark all that
	1	-	Improper medication administration, medication side effects, toxicity, anaphylaxis
	2	-	Injury caused by fall
	3	-	Respiratory infection (for example: pneumonia, bronchitis)
	4	-	Other respiratory problem
	5	-	Heart failure (for example: fluid overload)
	6	-	Cardiac dysrhythmia (irregular heartbeat)
	7	-	Myocardial infarction or chest pain
	8	-	Other heart disease
	9	-	Stroke (CVA) or TIA
	10	-	Hypo/Hyperglycemia, diabetes out of control
	11	-	GI bleeding, obstruction, constipation, impaction
	12	-	Dehydration, malnutrition
	13	-	Urinary tract infection
	14	-	IV catheter-related infection or complication
	15	-	Wound infection or deterioration
	16	-	Uncontrolled pain
	17	-	Acute mental/behavioral health problem
	18	-	Deep vein thrombosis, pulmonary embolus
	19	-	Scheduled treatment or procedure
	20	-	Other than above reasons
	UK	-	Reason unknown
(M0903)	Date		Last (Most Recent) Home Visit:
		'n	/// nonth / day / year
(M0906)	Disc		ge/Transfer/Death Date: Enter the date of the discharge, transfer, or death (at home) of the patient.
		'n	/// nonth / day / year